

A Covariant Road to Spatial Averaging in Cosmology : Scalar Corrections to the Cosmological Equations

Aseem Paranjape¹

*Tata Institute of Fundamental Research,
Homi Bhabha Road, Mumbai 400 005, India.*

This essay received an Honourable Mention in the Gravity Research Foundation's Essay Competition, 2007.

Abstract

A consistent approach to Cosmology requires an explicit averaging of the Einstein equations, to describe a homogeneous and isotropic geometry. Such an averaging will in general modify the Einstein equations. The averaging procedure due to Buchert has attracted considerable attention recently since it offers the tantalizing hope of explaining the phenomenon of dark energy through such corrections. This approach has been criticized, however, on the grounds that its effects may be gauge artifacts. We apply the fully covariant formalism of Zalaletdinov's Macroscopic Gravity and show that, after making some essential gauge choices, the Cosmological equations receive *spacetime scalar* corrections which are therefore observable in principle, and further, that the broad structure of these corrections is *identical* to those derived by Buchert.

¹e-mail address: aseem@tifr.res.in

Cosmology crucially assumes that the matter distribution in the Universe, when averaged on large enough scales, is homogeneous and isotropic. Assuming that the metric can also be “averaged” on these scales, one then models this averaged metric of spacetime as having the Friedmann-Lemaître-Robertson-Walker (FLRW) form with homogeneous and isotropic spatial sections, and applies Einstein’s General Relativity to determine the geometry of the Universe. The averaging operation is usually assumed implicitly and rather vaguely. It has long been known however [1], that any *explicit* averaging scheme for the metric g of spacetime and energy-momentum tensor T of matter, must necessarily yield corrections to the Einstein equations (which should ideally be imposed on length scales comparable to, say, the Solar System). This is a consequence of the Einstein tensor $E[g]$ being a nonlinear functional of the metric g : given some averaging operator $\langle \cdot \rangle$, in general one has $\langle E[g] \rangle \neq E[\langle g \rangle]$, and by using $E[\langle g \rangle] = \langle T \rangle$ as the field equations, one is ignoring these corrections.

Clearly, one needs a systematic averaging scheme within which to ask how large these corrections are. Among the averaging schemes available in the literature, only two – Buchert’s spatial averaging of scalars [2] and Zalaletdinov’s Macroscopic Gravity (MG) [3] – are capable of addressing the issue of averaging in General Relativity in a *nonperturbative* manner. (This is important since one expects the effects of averaging, if any, to show up only through nonlinear inhomogeneities.) Buchert’s approach has attracted considerable attention recently due partly to its tractability, but mainly because it offers the tantalizing hope of solving the dark energy problem *completely within the framework of classical General Relativity*. However, this approach has been criticized on the grounds that an averaging operation such as Buchert’s which is defined only for a particular $3+1$ splitting of spacetime, is likely to lead to observationally irrelevant gauge artifacts which could wrongly be interpreted as solving the dark energy problem.

While it is difficult to refute such criticism from within Buchert’s noncovariant ap-

proach, it might be possible to address the issue beginning with a *covariant* averaging scheme. Zalaletdinov’s MG is precisely such a scheme – fully covariant and mathematically elegant – which unfortunately comes with the somewhat steep price of extreme technical complexity. Nevertheless, the following question can be posed : Starting with the fully covariant structure of MG, and then making some appropriate gauge choices, is it possible to derive equations resembling Buchert’s modified FLRW equations? We answer this question in the affirmative [4], and address some of the issues that emerge in this construction. We begin by describing the broad structure of MG.

Zalaletdinov’s approach considers a general differentiable manifold $\mathcal{M}$ with metric g_{ab} , and defines a spacetime averaging operation for tensors which is then used to *construct* an “averaged” differentiable manifold $\bar{\mathcal{M}}$. The average of the affine connection on $\mathcal{M}$ is shown to itself behave like a connection, and is used as the affine connection for the abstract manifold $\bar{\mathcal{M}}$. The averaging operation is very sophisticated – it is defined using a Lie dragging of averaging regions along chosen vector fields, and ensures that the average of some tensor field $p_j^i(x)$, say, is itself a local tensor field $\langle p_j^i \rangle(x)$ on the manifold $\mathcal{M}$. One defines the (tensorial) connection correlation terms,

$$Z^a{}_{b[m]{}_{\underline{j}n}}{}^i = \left\langle \Gamma_{b[m]{}_{\underline{j}n}}^a \Gamma_{\underline{j}n}^i \right\rangle - \left\langle \Gamma_{b[m]}^a \right\rangle \left\langle \Gamma_{\underline{j}n}^i \right\rangle \quad ; \quad Z^a{}_{ijb} = 2Z^a{}_{ik}{}^k{}_{jb}, \quad (1)$$

where the square brackets denote antisymmetrization and the underlined indices are not antisymmetrized. It can then be shown that the averaged Einstein equations read

$$E_b^a = -\kappa T_b^a + C_b^a, \quad (2)$$

where $\kappa = 8\pi G_N$, E_b^a is the Einstein tensor for $\bar{\mathcal{M}}$, T_b^a is the averaged energy-momentum

tensor and the correlation tensor C_b^a is defined as

$$C_b^a = \left(Z_{ijb}^a - \frac{1}{2} \delta_b^a Z_{ijm}^m \right) G^{ij}, \quad (3)$$

where $G_{ab} = \langle g_{ab} \rangle$ is the metric on $\bar{\mathcal{M}}$, which we assume to be a highly symmetric space. The tensor C_b^a also satisfies $C_{b;a}^a = 0$, where the semicolon denotes covariant derivative on $\bar{\mathcal{M}}$.

We now make two important assumptions. Firstly, starting with some (formally unspecified) choice of $3 + 1$ splitting for the inhomogeneous spacetime, we consider the *time* averaging scale (denoted T) to be much smaller than the typical scale of time evolution of the metric. This is justified physically in the cosmological context since all measurements are performed over timescales much smaller than the Hubble expansion scale. Formally then, we consider the $T \rightarrow 0$, “spatial averaging limit” of MG. (This limit is intuitively seen to be well defined for averages of smooth functions, and this can also be verified explicitly.) Secondly, we assume that the averaging *length* scale (denoted L) is of order $\sim 150\text{-}200\text{Mpc}$, the scale at which cosmological homogeneity sets in. The averaged manifold $\bar{\mathcal{M}}$ is then assumed to admit a unit timelike vector field $\bar{v}^a$, which is orthogonal to spacelike hypersurfaces of constant curvature – in other words, $\bar{\mathcal{M}}$ is the FLRW spacetime. The spatial averaging limit is important since it ensures that the FLRW spacetime is a fixed point of the averaging scheme. Further treating the averaged energy-momentum tensor to have the form of a homogeneous and isotropic perfect fluid, one can write down two *scalar* equations – the (modified) Friedmann and Raychaudhuri equations – given by

$$\left(\frac{1}{a} \frac{da}{d\tau} \right)^2 = \frac{8\pi G_N}{3} \rho - \frac{1}{3} C_b^a \bar{v}^b \bar{v}_a \quad ; \quad \frac{1}{a} \frac{d^2 a}{d\tau^2} = -\frac{4\pi G_N}{3} (\rho + 3p) + \frac{1}{6} (C_a^a + 2C_b^a \bar{v}^b \bar{v}_a) . \quad (4)$$

We have assumed that the averaged metric takes the (flat, for simplicity) FLRW form in the synchronous gauge, with scale factor $a(\tau)$. The specific gauge chosen is irrelevant since these are scalar equations. ρ and p are the homogeneous energy density and pressure respectively as measured by “observers” in $\bar{\mathcal{M}}$ travelling on trajectories with tangent vector field $\bar{v}^a$. Physically these observers correspond to the *averaging domains* treated as physically infinitesimal cells – this is a refinement of the Weyl postulate which treated individual galaxies as travelling on hypersurface orthogonal trajectories. It is obvious from Eqn. (4) that the corrections to the standard cosmological equations arising from averaging of inhomogeneities, are *spacetime scalars*. In addition, we are forced to set the 3-vector $(C_A^0, A = 1, 2, 3)$ and traceless 3-tensor $(C_B^A - (1/3)\delta_B^A(C_J^J))$ parts of C_b^a to zero, due to the structure of E_b^a and the assumed form of T_b^a . This imposes additional consistency constraints on the inhomogeneous geometry.

The main question we wish to address concerns the *structure* of the corrections. Immediately, this leads to another question, namely, what choice of gauge for the *inhomogeneous* geometry, will lead to the averaged metric in the standard synchronous FLRW form? This question has no clear answer, and it is perhaps wiser to simply leave this gauge choice formally unspecified. It turns out however, that the form of the corrections simplifies if one begins with a *volume preserving gauge*, in which the metric determinant is constant. This happens because the structure of the MG formalism itself becomes very simple in the volume preserving gauge [3]. We will therefore display these results to highlight the broad structure of these corrections. Let us suppose that the inhomogeneous metric is written in coordinates wherein it takes the form

$$^{(\mathcal{M})}ds^2 = -\frac{dt^2}{h(t, \mathbf{x})} + h_{AB}(t, \mathbf{x})dx^A dx^B \quad ; \quad A, B = 1, 2, 3, \quad (5)$$

with $h = \det(h_{AB})$, so that $\sqrt{-g} = (1/\sqrt{h})(\sqrt{h}) = 1$. If we now assume that the “spatial”

average of h_{AB} is

$$\langle h_{AB} \rangle = \bar{a}^2(t) \delta_{AB}, \quad (6)$$

then it can be shown [4] that the full averaged metric must take the volume preserving form

$$^{(\bar{\mathcal{M}})}ds^2 = -\frac{dt^2}{\bar{a}^6(t)} + \bar{a}^2(t) \delta_{AB} dx^A dx^B, \quad (7)$$

which can be brought to synchronous form by the transformation $\tau = \int^t dt/\bar{a}^3$. Further, writing $a(\tau) = \bar{a}(t(\tau))$, the modified Cosmological equations now read

$$\left(\frac{1}{a} \frac{da}{d\tau} \right)^2 = \frac{8\pi G_N}{3} \rho - \frac{1}{6} [\mathcal{Q} + \mathcal{S}], \quad (8a)$$

$$\frac{1}{a} \frac{d^2 a}{d\tau^2} = -\frac{4\pi G_N}{3} (\rho + 3p) + \frac{1}{3} \mathcal{Q}, \quad (8b)$$

where $\mathcal{Q}$ and $\mathcal{S}$ are given by

$$\mathcal{Q} = \bar{a}^6 \left[\frac{2}{3} \left(\left\langle \frac{1}{h} \Theta^2 \right\rangle - \frac{1}{\bar{a}^6} ({}^F\Theta^2) \right) - 2 \left\langle \frac{1}{h} \sigma^2 \right\rangle \right], \quad (9a)$$

$$\mathcal{S} = \frac{1}{\bar{a}^2} \delta^{AB} \left[\left\langle ({}^3\Gamma_{AC}^J ({}^3\Gamma_{BJ}^C) \right\rangle - \left\langle \partial_A (\ln \sqrt{h}) \partial_B (\ln \sqrt{h}) \right\rangle \right]. \quad (9b)$$

Here Θ and σ^2 are the expansion and shear scalars respectively for the inhomogeneous geometry in the gauge (5), ${}^3\Gamma_{BC}^A$ is the Christoffel symbol constructed using h_{AB} , and the expansion scalar for the *averaged* geometry in the volume preserving gauge is given by $(1/\bar{a}^3)({}^F\Theta) = (3/\bar{a})(d\bar{a}/dt)$. $\mathcal{Q}$ and $\mathcal{S}$ are defined in the volume preserving gauge, but they are considered functions of the synchronous time τ in Eqns. (8), satisfying the

condition

$$(\partial_\tau \mathcal{Q} + 6\mathcal{Q}\partial_\tau(\ln a)) + (\partial_\tau \mathcal{S} + 2\mathcal{S}\partial_\tau(\ln a)) = 0, \quad (10)$$

which follows from $C_{b;a}^a = 0$. The additional consistency conditions mentioned earlier read

$$\delta^{JK} \left[\left\langle \sqrt{h} \Theta_{JB} {}^{(3)}\Gamma_{AK}^B \right\rangle - \left\langle \sqrt{h} \Theta_{JK} {}^{(3)}\Gamma_{AB}^B \right\rangle \right] = 0, \quad (11a)$$

$$\delta^{JK} \left\langle \frac{1}{\sqrt{h}} \Theta_K^B {}^{(3)}\Gamma_{JB}^A \right\rangle - \delta^{AJ} \left\langle \frac{1}{\sqrt{h}} \Theta {}^{(3)}\Gamma_{JB}^B \right\rangle = 0, \quad (11b)$$

$$\delta^{JK} \left\langle {}^{(3)}\Gamma_{JC}^A {}^{(3)}\Gamma_{KB}^C \right\rangle - \delta^{AJ} \left\langle {}^{(3)}\Gamma_{JC}^C {}^{(3)}\Gamma_{BK}^K \right\rangle = \frac{1}{3} \delta_B^A (\bar{a}^2 \mathcal{S}), \quad (11c)$$

with Θ_B^A the expansion tensor.

Our result contains some striking features. Firstly, despite the fact that the MG approach is very different from Buchert's spatial averaging (where, e.g., scalar equations are constructed *before* averaging as opposed to our Eqns. (4), see also Ref. [4]), our final results in Eqns. (8) and (10) are formally *identical* in structure to Buchert's modified Cosmological equations (see, e.g., Eqns. (10) and (13) in the first of Ref. [2]). We can therefore legitimately claim that corrections similar to those derived by Buchert arise as *spacetime scalars* in the covariant MG approach. Further, Eqns. (11) indicate that it is not sufficient to demand that an inhomogeneous metric average out to the FLRW form – additional correlations are also required to vanish and this imposes further conditions on the inhomogeneous metric.

To conclude, we have argued in the framework of Zalaletdinov's covariant MG, that the modifications to the standard FLRW equations are spacetime scalars, and hence in principle, observables. Moreover, under some simplifying assumptions we have demonstrated that the *structure* of these corrections, and the equations they satisfy, are formally *identical* to that of similar corrections derived by Buchert via a completely different

approach. The question of whether these corrections can account for the observed acceleration of FLRW models, while still open, is now on a concrete, observationally relevant footing.

References

- [1] G F R Ellis, in *General Relativity and Gravitation* (D. Reidel Publishing Co., Dordrecht), Eds. B. Bertotti *et al.*, (1984).
- [2] T Buchert, *Gen. Rel. and Grav.*, **32**, 105, (2000) [gr-qc/9906015]; *Gen. Rel. and Grav.*, **33**, 1381, (2001) [gr-qc/0102049].
- [3] R M Zalaletdinov, *Gen. Rel. & Grav.*, **25**, 673, (1993); *Gen. Rel. & Grav.*, **24**, 1015, (1992);
M Mars and R M Zalaletdinov, *J. Math. Phys.*, **38**, 4741, (1997) [dg-ga/9703002].
- [4] A Paranjape and T P Singh, [gr-qc/0703106], (2007), submitted.